
# Anti-Human CD11a (TP1/40)





#### PRODUCT DESCRIPTION

- Clone: TP1/40:
- Isotype: IgGl;
- Tested application: flow cytometry;
- Immunogen: The anti-CDIIa monoclonal antibody derives from PMA and Ionomycin activated T cells blast (Human);
- Species reactivity: Human;
- Storage instruction: store in the dark at 2-8 °C;
- Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN<sub>3</sub>);
- Recommended usage: Immunostep's CDIIa, clone TPI/40, is a monoclonal antibody intended for the identification and enumeration of human leucocytes using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using I test for I0<sup>6</sup> cells;
- Presentation: liquid;
- Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
- Purification: Affinity chromatography;
- Other names: Integrin alpha-L, CDII antigen-like family member A, Leukocyte adhesion glycoprotein LFA-I alpha chain, LFA-IA, Leukocyte function-associated molecule I alpha chain:
- Gene ID: 3683;
- Molecular weight: 170 180 kDa.

### 2. ANTIGEN DETAILS

**Large description:** The monoclonal antibody is directed against the CDIIa- antigen, located on the alpha-L chain of LFA-I complex (Lymphocyte Function-associated Antigen-I), which is expressed on mature immunocompetent lymphocytes and their neoplastic counterparts, granulocytes and monocytes<sup>(I-S)</sup>

## WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

### 4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

# REFERENCES

- Sanchez-Madrid F, Krensky AM, Ware CF, Robbins E, Strominger JL, Burakoff SJ, et al. Three distinct antigens associated with human T-lymphocyte-mediated cytolysis: LFA-1, LFA-2, and LFA-3. Proc Natl Acad Sci U S Al982 Dec;79(23):7489-93.
- Sanchez-Madrid F, Nagy JA, Robbins E, Simon P, Springer TA. A human leukocyte differentiation antigen family with distinct alpha-subunits and a common beta-subunit: the lymphocyte function-associated antigen (LFA-I), the C3bi complement receptor (OKMI/Mac-I), and the pI50,95 molecule. J Exp MedI983 Dec I;158(6):1785-803.
- Kishimoto TK, Hollander N, Roberts TM, Anderson DC, Springer TA. Heterogeneous mutations in the beta subunit common to the LFA-1, Mac-1, and p150,95 glycoproteins cause leukocyte adhesion deficiency. Celli987 Jul 17:50(2):193-202.
- Springer TA, Dustin ML, Kishimoto TK, Marlin SD. The lymphocyte function-associated LFA-1, CD2, and LFA-3 molecules: cell adhesion receptors of the immune system. Annu Rev Immunol1987:5:223-52.
- Knapp W. Leucocyte typing IV: white cell differentiation antigens. Oxford: Oxford University Press; 1989.

# 6. EXPLANATION OF SYMBOLS

	Fluorochrome
REF	Product reference
$\sum$	Content for <n> analysis</n>
	Regulatory Status
RUO	Research Use Only
•••	Manufacturer

# MANUFACTURED BY:

7.

# 444

#### IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C)

Campus de Unamuno 37007 Salamanca (Spain)

Telf./fax: (+34) 923 294 827
E-mail: info@immunostep.com

Revision N° 6 | Emission date: 03/02/202