# Anti-Human CD45 (D3/9)



	REF	Σ Σ	
FITC	45F1-100T	100 test	
PE	45PE1-100T	100 test	
PerCP	45PP1-100T	100 test	RUO
APC	45A1-100T	100 test	
CF-Blue	45CFB1-100T	100 test	

# 1. PRODUCT DESCRIPTION

- Clone: D3/9;
- Isotype: IgGl;
- Tested application: flow cytometry;
- Immunogen: The anti-CD45 monoclonal antibody derives from T cells from leukemic HPB-ALL;
- Species reactivity: Human;
- Storage instruction: store in the dark at 2-8 °C;
- Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN<sub>3</sub>);
- Recommended usage: Immunostep's CD45, clone D3/9 is a monoclonal antibody intended for the identification and enumeration of T200 protein using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 10<sup>6</sup> cells;
- Presentation: liquid;
- Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
- Purification: Affinity chromatography;
- Other names: Receptor-type tyrosine-protein phosphatase C, Leukocyte common antigen, L-CA, T200;
- Gene ID: 5788;
- Molecular weight: 180, 195, 205, 220 and 240 kDa.

# 2. ANTIGEN DETAILS

**Large description**: The monoclonal antibody is directed against the CD45-antigen, defined T200 or Leucocyte Common Antigen. The antibody reacts with all cells of the haemopoietic lineage, not with cells of other lineages.

CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases and associated with other surface antigen like CDI, CD2, CD3 and CD4.<sup>[1-4]</sup>

## 3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

### 4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

# REFERENCES

5.

- Krensky AM, Sanchez-Madrid F, Robbins E, Nagy JA, Springer TA, Burakoff SJ. The functional significance, distribution, and structure of LFA-I, LFA-2, and LFA-3: cell surface antigens associated with CTL-target interactions. J Immunol1983 Aug;131(2):611-6.
- Knapp W. Leucocyte typing IV : white cell differentiation antigens. Oxford: Oxford University Press; 1989.
- Marie-Cardine A, Maridonneau-Parini I, Fischer S. Activation and internalization of p56lck upon CD45 triggering of Jurkat cells. Eur J Immunol1994 Jun;24(6):1255-61.
- Escribano L, Orfao A, Villarrubia J, Diaz-Agustin B, Cervero C, Rios A, et al. Immunophenotypic characterization of human bone marrow mast cells. A flow cytometric study of normal and pathological bone marrow samples. Anal Cell Pathol1998;16(3):151-9.

### 6. EXPLANATION OF SYMBOLS

ι∎/ ∎	Fluorochrome
REF	Product reference
$\sum$	Content for <n> analysis</n>
	Regulatory Status
RUO	Research Use Only
	Manufacturer

7. MANUFACTURED BY: IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C) Campus de Unamuno 37007 Salamanca (Spain) Telf./fax: (+34) 923 294 827 E-mail: info@immunostep.com www.immunostep.com