Diluent Buffer (1X – Ready to use)

Sample diluent that minimizes non-specific binding, cross-reactivity and matrix interference (without indigo carmine E-135).

1. INTRODUCTION


2. INSTRUCTIONS FOR USE

.

4.

Before use, temper the buffer between +18°C y +24°C (room temperature) for 30 minutes.

 Samples are diluted 1:20 sample diluent buffer (1X) and mixed with a cortex agitator according the below figure. Diluted samples should be assayed within 8 hours.



3. RELATED PRODUCTS

- Covid Multiplex (Ref. IMS0509)
- Covid ELISA (Refs. IMS2905, IMS2906 and IMS2907)

IMMUNOSTEP S.L.

E-mail:

Telf./fax: (+34) 923 294 827

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C)

Campus de Unamuno 37007 Salamanca (Spain)

info@immunostep.com

www.immunostep.com

ExoStep ELISA (Ref. EXO2506 and EXO2508)

MANUFACTURED BY

Reference: IMS

Volume: 10 mL

Storage: 2-8°C

pH-value: 7.2 ± 0.2

Preservative: contains < 0.0014 % [w/w] reaction mass of CMIT/MIT (3:1)

Buffer Dilue

lin (+34) 923 29 48 27 1

S0514

(10mL)

Expiry date when stored please refer to the label on the bottle. unopened:

FOR RESEARCH USE ONLY, NOT FOR DIAGNOSTIC USE.

Immunostep