




Anti-Human CD33 (HIM3-4)

	REF	Σ		
FITC	33F-100T	100 test	20 μ L/test	
PE	33PE-100T	100 test	20 μ L/test	
APC	33A-100T	100 test	20 μ L/test	

1. PRODUCT DESCRIPTION

Clone: HIM3-4;
Isotype: IgG1;
Tested application: flow cytometry;
Immunogen: The anti-CD33 monoclonal antibody derives from KG1a Cell Line;
Species reactivity: Human, Cross-Reactivity: Chimpanzee;
Storage instruction: store in the dark at 2-8 °C;
Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₃);
Recommended usage: Immunostep's CD33, clone HIM3-4 is a monoclonal antibody intended for the identification and enumeration of Siglec-3 using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 10⁶ cells;
Presentation: liquid;
Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
Purification: Affinity chromatography;
Other names: Myeloid cell surface antigen CD33, Sialic acid-binding Ig-like lectin 3, Siglec-3, gp67, p67;
Gene ID: 945;
Molecular weight: 67 rDa.

2. ANTIGEN DETAILS

Large description: The monoclonal antibody is directed against the CD33-antigen, which is expressed on human myelomonocytic cells; monocytes, granulocytes (weakly), myeloid progenitors and mast cells. The monoclonal antibody reacts in the bone marrow from myeloblasts up to myelocytes. CD33-antigen is found on CFU-GEMM, CFU-GM, CFU-G, CFU-M and with erythroid CFU-E but not on earlier precursors. CD33 does not react with normal human peripheral B-cells, T-cells and platelets. CD33 reacts weakly with blast cells in 70% of patients with Acute Myeloid Leukaemia (AML) and in 30% of adult patients with Acute Lymphoblastic Leukaemia (ALL).⁽¹⁻⁶⁾

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.








Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

5. REFERENCES

- Favaloro EJ, Bradstock KF, Kabral A, Grimsley P, Zowtyj H, Zola H. Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. Br J Haematol1988 Jun;69(2):163-71.
- Favaloro EJ, Moraitis N, Koultis J, Exner T, Bradstock KF. Endothelial cells and normal circulating haemopoietic cells share a number of surface antigens. Thromb Haemost1989 Apr 25;61(2):217-24.
- Freeman SD, Kelm S, Barber EK, Crocker PR. Characterization of CD33 as a new member of the sialoadhesin family of cellular interaction molecules. Blood1995 Apr 15;85(8):2005-12.
- Nakamura Y, Noma M, Kidokoro M, Kobayashi N, Takei M, Kurashima S, et al. Expression of CD33 antigen on normal human activated T lymphocytes. Blood1994 Mar 1;83(5):1442-3.
- Ulyanova T, Blasioli J, Woodford-Thomas TA, Thomas ML. The sialoadhesin CD33 is a myeloid-specific inhibitory receptor. Eur J Immunol1999 Nov;29(11):3440-9.
- Knapp W. Leucocyte typing IV : white cell differentiation antigens. Oxford: Oxford University Press; 1989.

6. EXPLANATION OF SYMBOLS

	Form
	Catalog reference
	Contains sufficient for <n> test
	Quantity per test
	Regulatory Status
	Research Use Only
	Manufacturer

7. MANUFACTURED BY:



IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n
 Cancer Research Center (C.I.C)
 Campus de Unamuno
 37007 Salamanca (Spain)
Telf./fax: (+34) 923 294 827
E-mail: info@immunostep.com
www.immunostep.com