Anti-Human CD36 (CLB-IVC7)













FITC 36F2-100T 100 test 20 µL/test
PerCP-Cyanine 5.5 36PP5.52-100T 100 test 5 µL/test
CF-Blue 36CFB2-100T 100 test 5 µL/test



PRODUCT DESCRIPTION

Clone: CLB-IVC7;

Isotype: IgGl;

Tested application: flow cytometry;

Immunogen: The CD36 antibody, clone CLB-IVC7, is derived from the hybridization of Sp2/O mouse myeloma cells with spleen cells isolated from BALB/c mice immunized with human monocytes:

Species reactivity: Human;

Storage instruction: store in the dark at 2-8 °C;

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₃);

Recommended usage: Immunostep's CD36, cloneCLB-IVC7, is a monoclonal antibody intended for the identification and enumeration of platelets, macrophages, endothelial cells, early erythroid cells and megakaryocytes using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 10⁶ cells;

Presentation: liquid;

Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;

Purification: Affinity chromatography;

Other names: GPIV, GP4, GP3B, thrombospondin receptor, PASIV, FAT, and SCARB3;

Gene ID: 948;

Molecular weight: 88 kDa.

2. ANTIGEN DETAILS

Large description: CD36 is a transmembrane, highly glycosylated, glycoprotein expressed by monocytes, macrophages, platelets, microvascular endothelial cells, early erythroid cells and megakaryocytes and adipose tissues and weak with B cells. Detection of CD36 on human blood T cells is spurious, due mainly to interaction with contaminating platelets which express CD36 robustly5.

Platelet glycoprotein IV (GP IV)(GPIIIb) (CD36 antigen) is also called GPIV, OKM5-antigen or PASIV. CD36 recognises oxidized low density lipoprotein, long chain fatty acids, anionic phospholipids, collagen types I, IV and V, thrombospholipid (TSP) and Plasmodium falciparum infected erythrocytes. Binds long chain fatty acids and may function in the transport and/or as a regulator of fatty acid transport. $^{(i-4)}$

WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

Revision N° 7 | Emission date: 09/02/2021

REFERENCES

5.

- Alessio M, Ghigo D, Garbarino G, Geuna M, Malavasi F. Analysis of the human CD36 leucocyte differentiation antigen by means of the monoclonal antibody NLO7. Cell lmmunoll991 Oct 15;137(2):487-500.
- Alessio M, Roggero S, Bussolino F, Saitta M, Malavasi F. Characterization of the murine monoclonal antibody NLO7 specific for the human thrombospondin receptor (CD36 molecule). Curr Stud Hematol Blood Transfus1991/58):182-6.
- Alessio M, Greco NJ, Primo L, Ghigo D, Bosia A, Tandon NN, et al. Platelet activation and inhibition of malarial cytoadherence by the anti-CD36 IgM monoclonal antibody NLO7. Blood1993 Dec 15:82(12):3637-47.
- Gruarin P, Ulliers D, Thorne RF, Alessio M. Methionine 156 in the immunodominant domain of CD36 contributes to define the epitope recognized by the NL07 MoAb. Mol Cell Biochem2000 Nov;214(1-2):89-95.
- Zamora C, Cantó E, Nieto JC, Ortiz MA, Diaz-Torné C, Diaz-Lopez C, Llobet JM, Juarez C, Vidal S. Functional consequences of platelet binding to T lymphocytes in inflammation. J Leukoc Biol. 2013 Sep;94(3):521-9. doi: 10.1189/jlb.0213074. Epub 2013 Jun 25. PMID: 23801652.

6. EXPLANATION OF SYMBOLS

\• /	Form
REF	Catalog reference
Σ	Contains sufficient for <n> test</n>
\Diamond	Quantity per test
	Regulatory Status
RUO	Research Use Only
•••	Manufacturer

MANUFACTURED BY:

7.



IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C)

> Campus de Unamuno 37007 Salamanca (Spain)

Telf./fax: (+34) 923 294 827 E-mail: info@immunostep.com

www.immunostep.com