Anti-Human CD43 (TP1/36)



	REF	\sum	\Diamond	
PURE	43PU	1 mg	1 mg/ml	1
FITC	43F-100T	100 test	20 μL/test	
PE	43PE-100T	100 test	20 μL/test	
PerCP	43PP-100T	100 test	5 μL/test	RUO
PerCP-Cyanine5.5	43PP5.5-100T	100 test	5 µL/test	
APC	43A-100T	100 test	20 µL/test	
APC-C750	43AC750-100T	100 test	5 μL/test	
CF-Blue	43CFB-100T	100 test	5 μL/test	

PRODUCT DESCRIPTION

Clone: TP1/36; Isotype: IqG1;

Tested application: flow cytometry;

Immunogen: The anti-CD43 monoclonal antibody derives from PMA and lonomycin

activated T cells blast (Human);

Species reactivity: Human;

Storage instruction: store in the dark at 2-8 °C;

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaNa);

Recommended usage: Immunostep's CD43, clone TP1/36 is a monoclonal antibody intended for the identification and enumeration of thymocytes and T lymphocytes using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 106 cells;

Presentation: liquid:

Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;

Purification: Affinity chromatography;

Other names: Leukosialin, Galactoglycoprotein, GALGP, Leukocyte sialoglycoprotein, Sialophorin, GPL115, LSN, Ly-48, Leukosialin, qp115;

Gene ID: 6693:

Molecular weight: 95 - 135 kDa.

2. ANTIGEN DETAILS

Large description: Reacts with a single-pass type I protein found on cell surface of thymocytes. T-lymphcytes, neutrophis, plasma cells and myelomas but is not present on most peripheral blood B cells. CD43 has a high content of sialic acid and O-linked carbohydrate structures. (1-4)

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

REFERENCES

5.

- Borche L. Lozano F. Vilella R. Vives J. CD43 monoclonal antibodies recognize the large sialoglycoprotein of human leukocytes. Eur J Immunol1987 Oct;17(10):1523-6.
- Carlsson SR, Fukuda M. Isolation and characterization of leukosialin, a major sialoglycoprotein on human leukocytes. J Biol Chem1986 Sep 25;261(27):12779-86.
- Knapp W. Leucocyte typing IV: white cell differentiation antigens. Oxford: Oxford University Press; 1989.
- Remold-O'Donnell E, Zimmerman C, Kenney D, Rosen FS. Expression on blood cells of sialophorin, the surface glycoprotein that is defective in Wiskott-Aldrich syndrome. Blood1987 Jul:70(1):104-9.

6. **EXPLANATION OF SYMBOLS**

\ . }	Form
REF	Catalog reference
\sum	Contains sufficient for <n> test</n>
\Diamond	Quantity per test
	Regulatory Status
RUO	Research Use Only
***	Manufacturer

7. MANUFACTURED BY:



IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C)

Campus de Unamuno 37007 Salamanca (Spain)

Telf./fax: (+34) 923 294 827 E-mail: info@immunostep.com

www.immunostep.com