

Lambda Light Chain (F'ab2 Polyclonal)



PURE	LAMBDAF3PU	1 mg	1 mg/ml	RUO
FITC	LF2-100T	100 test	20 µL/test	
PE	LPE2-100T	100 test	20 µL/test	
APC	LA2-100T	100 test	10 µL/test	
APC-C750	LAC7502-100T	100 test	5 µL/test	

1. PRODUCT DESCRIPTION

Clone: Polyclonal;
Isotype: Rabbit F(ab')₂ IgG;
Tested application: flow cytometry;
Immunogen: Polyclonal immunoglobulin light chains of lambda type isolated from a pool of human sera for Rabbit Anti-Human Lambda Light Chains;
Species reactivity: Human;
Storage instruction: store in the dark at 2-8 °C;
Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₃);
Recommended usage: Immunostep's lambda F'ab 2, is intended for simultaneous detection and enumeration of B lymphocytes bearing lambda light chains in peripheral blood using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 10 µl/10⁶ cells;
Presentation: liquid;
Source: Supernatant proceeding from sera for Rabbit Anti-Human Kappa Light Chains;
Purification: Affinity chromatography;
Other names: T Cell Receptor beta; TCRB; TRB; TRB@; TCR VB3-CBI;
Gene ID: 28639.
Molecular weight: 34 kDa.

2. ANTIGEN DETAILS

Large description: The evaluation of cell surface Kappa/Lambda expression can identify clonally restricted B lymphocyte populations and thus can aide in the diagnosis of hematologic malignancy. Several B cell disorders are associated with decreased levels of Kappa/Lambda at the cell surface⁽¹⁻⁴⁾.

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.








Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

5. REFERENCES

- Gandini D, Lanza F, Latorraca A, Levato F, Del Senno L, Castoldi G. Immunophenotypic and genotypic characterization of B-cell chronic lymphocytic leukemia patients from northern Italy. *Haematologica*1993 Jan-Feb;78(1):18-24.
- Johnson A, Olofsson T. Flow cytometric clonal excess analysis of peripheral blood, routine handling, and pitfalls in interpretation. *Cytometry*1993;14(2):188-95.
- Cartron G, Linassier C, Bremond JL, Desablens B, Georget MT, Fimbel B, et al. CD5 negative B-cell chronic lymphocytic leukemia: clinical and biological features of 42 cases. *Leuk Lymphoma*1998 Sep;31(1-2):209-16.
- Braylan RC, Orfao A, Borowitz MJ, Davis BH. Optimal number of reagents required to evaluate hematolymphoid neoplasias: results of an international consensus meeting. *Cytometry*2001 Feb 15;46(1):23-7.

6. EXPLANATION OF SYMBOLS

	Form
	Catalog reference
	Contains sufficient for <n> test
	Quantity per test
	Regulatory Status
	Research Use Only
	Manufacturer

7.

MANUFACTURED BY:



IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n
 Cancer Research Center (C.I.C)
 Campus de Unamuno
 37007 Salamanca (Spain)

Tel./fax: (+34) 923 294 827

E-mail: info@immunostep.com
www.immunostep.com