Multicolor cocktails TBNK





PerCp-Cyanine5.5/ APC

1. PRODUCT DESCRIPTION

Tested application: flow cytometry

Species reactivity: Human

Storage instruction: store in the dark at 2-8 °C

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN 3).

Recommended usage: TBNK reagent panel, which includes the combinations CD3 FITC / CD4 PE-Cyanine7 / CD8 APC-Cyanine7 / (CD16+CD56) PE / CD45 PerCP-Cy55 / CD19 APC and CD3 FITC / (CD16+CD56) PE / CD45 PerCP / CD19 APC, is intended for in vitro diagnostic use for the qualitative identification of major lymphocyte subsets — T cells (CD3, CD4, CD8), B cells (CD19, and NK cells (CD16+CD56) — in human peripheral blood samples by flow cytometry.

This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using I test for IO 6 cells.

Presentation: liquid

Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma. Purification: Affinity chromatography.

Reagent Composition TBNK1

ANTIBODY	FLUOROCHROME	CLONE	ISOTYPE
CD3	- FITC	33-2A3	lgG2a
CD16	PE	3G8	lgG1
CD56	PE	B-A19	lgG1
CD45	PerCP	D3/9	lgG1
CD19	APC	A3-B1	lgG2a

Reagent Composition TBNK2

ANTIBODY	FLUOROCHROME	CLONE	ISOTYPE
CD3	FITC	33-2A3	IgG2a
CD16	PE	3G8	-lgG1
CD56	-PE	B-A19	-lgG1
CD4	PE/Cyanine7	SIK3	-lgGl
CD45	-PerCP/Cyanine5.5	D3/9	-lgG1
CD19	APC	A3-B1	-lgG2a
CD8	APC/C750TM	- 143-44	lgG1

2. ANTIGEN DETAILS

CD3 (clone 33-2A3) recognizes the ε-chain of the CD3/TCR complex, which is expressed on all mature T lymphocytes. This antigen is a defining marker of the T-cell lineage and is essential for Tcell receptor signaling. CD3 expression is absent on B cells, NK cells, and other leukocyte subsets, allowing for specific identification of T cells by flow cytometry.

CD4 (clone SK3) identifies a glycoprotein expressed predominantly on helper/inducer T lymphocytes. CD4 acts as a co-receptor for MHC class II molecules and plays a central role in T-cell activation and immune regulation. It is also found at low levels on monocytes.

CD8 (clone 143-44) recognizes a cell surface glycoprotein that interacts with MHC class I molecules. It is expressed mainly on cytotoxic/suppressor T lymphocytes and a subset of NK cells. CD8 enhances adhesion to target cells and contributes to T-cell activation.

CDI6 (clone 3G8) binds to the low-affinity Fc γ RIII receptor for IgG, which is present on NK cells, neutrophils, and some monocyte subsets. It is widely used to identify NK cells in combination with CD56, although it can also label granulocytes.

CD56 (clone B-A19) targets the neural cell adhesion molecule (NCAM), which is expressed on natural killer (NK) cells and a subset of T lymphocytes (CD3 $^+$ CD56 $^+$ NKT-like cells). CD56 is essential for the identification of NK populations in peripheral blood.

CD45 (clone D3/9) binds to all isoforms of the leukocyte common antigen (LCA), which is present on all human leukocytes, including lymphocytes, monocytes, granulocytes, eosinophils, and basophils. CD45 is used to distinguish leukocytes from non-hematopoietic cells and debris.

CDI9 (clone A3-BI) recognizes a transmembrane protein expressed throughout B cell development, excluding plasma cells. It is a pan-B cell marker that does not react with T cells, monocytes, or granulocytes, allowing for specific detection of B lymphocytes.

Please, refer to www.immunostep.com technical support for more information.

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labeling at the time of delivery to the customer. Immunostep disclaims hereby other warranties. Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. REFERENCES

- de Vries E, Koene HR, Vossen JM, Gratama JW, von dem Borne AE, Waaijer JL, Haraldsson A, de Haas M, van Tol MJ. Identification of an unusual Fc gamma receptor Illa (CDI6) on natural killer cells in a patient with recurrent infections. Blood. 1996 Oct 15
- Blumberg RS, Ley S, Sancho J, Lonberg N, Lacy E, McDermott F, Schad V, Greenstein JL, Terhorst C. Structure of the T-cell antigen receptor: evidence for two CD3 epsilon subunits in the T-cell receptor-CD3 complex. Proc Natl Acad Sci U S A. 1990 Sep;87(18)
- Guan F, Wang X, He F. Promotion of cell migration by neural cell adhesion molecule (NCAM) is enhanced by PSA in a polysialyltransferase-specific manner. PLoS One. 2015 Apr 17
- Guan F, Wang X, He F. Promotion of cell migration by neural cell adhesion molecule (NCAM) is enhanced by PSA in a polysialyltransferase-specific manner. PLoS One. 2015 Apr 17

5. EXPLANATION OF SYMBOLS

 Form

 REF
 Catalog reference

 V
 Contains sufficient for <n> test

 Quantity per test
 Image: Contains sufficient for <n> test

 Regulatory Status
 Research Use Only

 Research Use Only
 Manufacturer

MANUFACTURED BY:

6.



Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C) Campus de Unamuno 37007 Salamanca (Spain) Telf./fax: (+34) 923 294 827 E-mail: info@immunostep.com www.immunostep.com