








Anti-Human CD117 (104D2)

					
PE	I17PE-100T	100 test	20 µL/test	0,05 mg/ml	
APC	I17A-100T	100 test	20 µL/test	0,05 mg/ml	
PerCP-Cyanine5.5	I17PPC5.5-100T	100 test	5 µL/test	0,2 mg/ml	
PE-Cyanine7	I17PC7-100T	100 test	5 µL/test	0,2 mg/ml	

1. PRODUCT DESCRIPTION

Clone: 104D2;
Isotype: IgG1;
Tested application: flow cytometry;
Immunogen: The anti-CD117 monoclonal antibody derives from MOLM-1 megakaryocytic cell line;
Species reactivity: Human, Cross-Reactivity: Cynomolgus, Cattle (Bovine, Cow);
Storage instruction: store in the dark at 2-8 °C;
Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₃);
Recommended usage: Immunostep's CD117, clone 104D2, is a monoclonal antibody intended for the identification and enumeration of lymphoid progenitor using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 10⁶ cells;
Presentation: liquid;
Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
Purification: Affinity chromatography;
Other names: Stem cell factor receptor, c-kit, mast cell growth factor receptor, steel factor receptor;
Gene ID: 3815;
Molecular weight: 145 kDa.

2. ANTIGEN DETAILS

Large description: This antibody reacts with the CD117 antigen, which is expressed within the haematopoietic compartment on approximately 50 % of CD34+ progenitors engaged in erythrocytic, myelo-monocytic and megakaryocytic differentiation. The 104D2 monoclonal antibody reacts with human CD117, also known as c-Kit, Steel factor receptor and stem cell factor receptor. A member of the tyrosine kinase receptor family, this 145 kDa molecule is expressed by hematopoietic progenitor cell subsets and mast cells. The interaction of c-Kit and Steel factor promotes proliferation and differentiation of hematopoietic progenitor cells and mast cell differentiation.⁽¹⁻⁵⁾

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.









Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

REFERENCES

- Gonzalez-de-Olano D, Matito A, Orfao A, Escribano L. Advances in the understanding and clinical management of mastocytosis and clonal mast cell activation syndromes. F1000Res;5:2666.
- Columbo M, Horowitz EM, Botana LM, MacGlashan DW, Jr., Bochner BS, Gillis S, et al. The human recombinant c-kit receptor ligand, rhSCF, induces mediator release from human cutaneous mast cells and enhances IgE-dependent mediator release from both skin mast cells and peripheral blood basophils. J Immunol1992 Jul 15;149(2):599-608.
- Ikuta K, Uchida N, Friedman J, Weissman IL. Lymphocyte development from stem cells. Annu Rev Immunol1992;10:759-83.
- Werfel T, Zwirner J, Oppermann M, Sieber A, Begemann G, Drommer W, et al. CD88 antibodies specifically bind to C5aR on dermal CD117+ and CD14+ cells and react with a desmosomal antigen in human skin. J Immunol1996 Aug 15;157(4):1729-35.
- Bravo P, Agustin BD, Bellas C, Gonzalez D, Camara C, Fuertes IF, et al. Expression of high amounts of the CD117 molecule in a case of B-cell non-Hodgkin's lymphoma carrying the t(14;18) translocation. Am J Hematol2000 Apr;63(4):226-9.

EXPLANATION OF SYMBOLS

	Form
	Catalog reference
	Contains sufficient for > test
	Quantity per test
	Regulatory Status
	Research Use Only
	Concentration
	Manufacturer

MANUFACTURED BY:



IMMUNOSTEP S.L.
Address: Avda. Universidad de Coimbra, s/n
 Cancer Research Center (C.I.C)
 Campus de Unamuno
 37007 Salamanca (Spain)
 (+34) 923 294 827
Telf./fax:
E-mail: info@immunostep.com
www.immunostep.com