Anti-Human CD1a (HI149)















0,05 mg/ml 20 uL/test 20 uL/test 0.05 ma/ml



1AA-100T 100 test 1. PRODUCT DESCRIPTION

1AF-100T 100 test

Clone: HI149:

Isotype: IgG1;

Tested application: flow cytometry;

Immunogen: The anti-CDIa monoclonal antibody derives from leukemia cells;

Species reactivity: Human;

Storage instruction: store in the dark at 2-8 °C;

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09%

Recommended usage: Immunostep's CDIa, clone HI149^(1, 2), is a monoclonal antibody intended for the identification and enumeration of human thymocyte antigen, present thymocytes using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using I test for 106 cells;

Presentation: liquid;

Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;

Purification: Affinity chromatography;

Other names: Leu-6, T6, R4;

Gene ID: 909:

Molecular weight: Iq superfamily, MHC I-like molecule, type I transmembrane glycoprotein, 49 kDa.

2. ANTIGEN DETAILS

Large description: This antibody reacts with the CDIa antigen which is found on human thymocytes and certain thymomas, on Langerhans cells in the skin, dermal dendritic cells and brain astrocytes. The antigen is a membrane glycoprotein which is expressed non-covalently with b2 - microglobulin and is closely similar to human thymocyte antigen-1. (3-5)

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

5. **REFERENCES**

- Knapp W. Leucocyte typing IV: white cell differentiation antigens, Oxford: Oxford University Press: 1989.
- Schlossman SF. Leucocyte typing V: white cell differentiation antigens: proceedings of the Fifth International Workshop and Conference: held in Boston, USA, 3-7 November, 1993. Oxford: Oxford University Press; 1995.
- Calabi F, Bradbury A. The CDI system. Tissue Antigens1991 Jan;37(1):1-9.
- Escribano L, Orfao A, Villarrubia J, Diaz-Agustin B, Cervero C, Rios A, et al. Immunophenotypic characterization of human bone marrow mast cells. A flow cytometric study of normal and pathological bone marrow samples. Anal Cell Pathol1998;16(3):151-9.
- Hanau D, Schmitt DA, Bieber T, Schmitt D, Cazenave JP. Possible mechanism of action of CDIa antigens. J Invest Dermatol1990 Nov:95(5):503-5.

6. **EXPLANATION OF SYMBOLS**

\.	Form
REF	Catalog reference
\sum	Contains sufficient for > test
\Diamond	Quantity per test
	Regulatory Status
RUO	Research Use Only
[A]	Concentration
	Manufacturer

7. MANUFACTURED BY:



Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C) Campus de Unamuno 37007 Salamanca (Spain) Telf./fax: (+34) 923 294 827 info@immunostep.com www.immunostep.com