

Anti-Human CD38 (HB-7)


REF

RUO

APC-C750

38AC752-100T

100 test

1. PRODUCT DESCRIPTION

Clone: HB7

Isotype: Mouse IgG1, k

Tested application: flow cytometry

Immunogen: The anti-CD38 monoclonal antibody derives from BJAB human B cell line.

Species reactivity: Human

Storage instruction: store in the dark at 2-8 °C

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN3).

Recommended usage: Immunostep's CD38, clone HB7, is a monoclonal antibody intended for the identification and enumeration of plasma cells using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 10 6 cells.

Presentation: liquid

Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma.

Purification: Affinity chromatography.

2. ANTIGEN DETAILS

Large description: The HB-7 monoclonal antibody reacts with the human CD38 molecule, a type II transmembrane protein. Expression of CD38 is bimodal during B cell development, modulating from high in immature cells to low in intermediate ones and back to high on mature B cells. Additionally CD38 is found in a variety of tissues and other hematopoietic cell. The CD38 antigen acts as a bifunctional ectoenzyme that catalyzes both the synthesis and the hydrolysis of a Ca++ mobilizing agent, cyclic ADP-ribose. This intracellular calcium plays an important role in cell signaling pathways. (1-4)

Other Names: TIO, ADPribosyl cyclase, and cyclic ADP ribose hydrolase I.

Gene ID: 952

Molecular weight: 45 kDa

For research use only, not for diagnostic procedures. Please, refer to www.immunostep.com technical support for more information.

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. REFERENCES

1. Reinherz EL, Kung PC, Goldstein G, Levey RH, Schlossman SF. Discrete stages of human intrathymic differentiation: analysis of normal thymocytes and leukemic lymphoblasts of T-cell lineage. *Proc Natl Acad Sci U S A* 1980 Mar;77(3):1588-92.
2. Knapp W. Leucocyte typing IV : white cell differentiation antigens. Oxford: Oxford University Press; 1989.
3. Jackson DG, Bell JI. Isolation of a cDNA encoding the human CD38 (TIO) molecule, a cell surface glycoprotein with an unusual discontinuous pattern of expression during lymphocyte differentiation. *J Immunol* 1990 Apr 1;144(7):2811-5.
4. Terstappen LW, Huang S, Safford M, Lansdorp PM, Loken MR. Sequential generations of hematopoietic colonies derived from single nonlineage-committed CD34+CD38- progenitor cells. *Blood* 1991 Mar 15;77(6):1218-27.

5. EXPLANATION OF SYMBOLS



Form



Catalog reference



Contains sufficient for > test



Regulatory Status



Quantity per test



Research Use Only



Concentration



Manufacturer

6. MANUFACTURED BY:

IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n
Cancer Research Center (C.I.C)
Campus de Unamuno
37007 Salamanca (Spain)
Telf./fax: (+34) 923 294 827
E-mail: info@immunostep.com
www.immunostep.com