

Anti-Human CD2 (RPA-2.10)



REF



StarBright™ Dye UltraViolet 445

25BUV4452-100T

100 test

Intended use for StarBright™ conjugates

1. PRODUCT DESCRIPTION

Clone: RPA-2.10

Isotype: Mouse IgG1, κ

Tested application: Flow cytometry

Species reactivity: Human

Storage instruction: Store in the dark at 2-8 °C

Storage buffer: Aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN³).

Field of Use for StarBright™ Dye conjugates: for use in in-house developed assays (IHA) or laboratory-developed tests (LDT).

Recommended usage: Immunostep's CD2, clone RPA-2.10, is a monoclonal antibody intended for the identification and enumeration of the CD2 antigen, a 50 kDa type I transmembrane glycoprotein also known as LFA-2, TII, and sheep red blood cell receptor (SRBC-R). This member of the immunoglobulin superfamily is expressed on thymocytes, T lymphocytes, natural killer (NK) cells, and subsets of thymic B cells. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using I test for 10⁶ cells.

Presentation: liquid

Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma.

Purification: Affinity chromatography.

2. ANTIGEN DETAILS

Large description: This antibody reacts with the CD2 antigen (also known as LFA-2, TII, or SRBC-R), a monomeric type I transmembrane glycoprotein of 45–58 kDa, expressed on all peripheral blood T lymphocytes, more than 95% of thymocytes, and a subset of natural killer (NK) cells. CD2 is one of the earliest markers of the T-cell lineage and is not expressed on B lymphocytes¹

CD2 plays a central role in T cell adhesion and activation. It interacts primarily with CD58 (LFA-3), a ligand that stabilizes the contact between T cells and antigen-presenting or target cells. Although the CD2–CD58 interaction has relatively low affinity in solution, it is significantly strengthened at the two-dimensional interface of cell-cell contact, especially upon T cell activation. This process involves increased CD2 expression, enhanced lateral mobility, and eventual fixation of CD2–CD58 complexes at the immunological synapse, promoting sustained intercellular adhesion and signaling^{2,3}.

Functionally, monoclonal antibodies against CD2, such as RPA-2.10, inhibit rosette formation with sheep erythrocytes, confirming CD2's role as or in association with the erythrocyte receptor. RPA-2.10 also blocks mixed lymphocyte reactions, further demonstrating its involvement in immune modulation⁴.

CD2 deficiency has been associated with reduced intestinal inflammation and altered immune responses. Dysregulation of CD2 expression or function has been implicated in conditions such as immune deficiency due to thymic absence and penile squamous cell carcinoma. The RPA-2.10 clone cross-reacts with non-human primates and pigs, making it suitable for comparative immunological studies².

Other Names: LFA-2, TII, SRBC-R

Gene ID: 914

Molecular weight: 50 kDa

Revision N° 1 | Emission date: 01/07/2025

3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties. Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use. Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

5. REFERENCES

- Shibata Y, Nakamura H, Kato S, Tomoike H. Cellular detachment and deformation induce IL-8 gene expression in human bronchial epithelial cells. *J Immunol.* 1996 Jan 15;156(2):772-7. PMID: 8543832.
- O'Rourke AM, Ybarrondo B, Mescher MF. CD8 and antigen-specific T cell adhesion cascades. *Semin Immunol.* 1993 Aug;5(4):263-70. doi: 10.1006/smim.1993.1030. PMID: 8219104.
- Davis SJ, van der Merwe PA. CD2: an exception to the immunoglobulin superfamily concept? *Science.* 1996 Aug 30;273(5279):1241-2. doi: 10.1126/science.273.5279.1241. PMID: 8703062.
- Monoclonal antibodies specific for the erythrocyte binding domain of the T cell adhesion molecule CD2 inhibit T cell rosette formation. *J Immunol.* 148(3):896-903.

6. EXPLANATION OF SYMBOLS



Form

REF

Catalog reference



Contains sufficient for <n> test



Regulatory Status



Quantity per test

RUO

Research Use Only



Manufacturer

7. MANUFACTURED BY: IMMUNOSTEP S.L.



Address: Avda. Universidad de Coimbra, s/n
Cancer Research Center (C.I.C)
Campus de Unamuno
37007 Salamanca (Spain)

Telf./fax: (+34) 923 294 827

E-mail: info@immunostep.com

www.immunostep.com